

***The status of selected next
generation double-beta decay
experiments***


Frank Avignone

University of South Carolina

7th Patras Workshop on axions, WIMPS and WISPS

Mykonos, Greece June 26th-July 1st, 2011

○ ● ● The Neutrino Hierarchy Problem



The neutrino oscillation experiments analyzed all together give us this picture; however, there are only bounds on the lightest neutrino mass eigenstate!



**Double-
beta decay**

0⁺ Te-130

¹³⁰₅₂Te

ββ

5⁺ I-130

¹³⁰₅₃I

β


Effects of pairing forces

**Weak decay to I-130
energetically forbidden**

2527.518 (13) keV FSU

$Q_{\beta\beta} = 2528 \pm 1.3 \text{ keV}$

Xe-130






Motivation for $0\nu\beta\beta$ Search

$${}^Z A \Rightarrow {}^{Z+2} A + 2e^{-}$$

- Immediate Implications of Discovery:
 - Neutrino is Majorana (own antiparticle)
 - Total Lepton Number is not conserved
 - Neutrino has mass (known)
- Well-studied example: Exchange of virtual neutrino

$2\nu\beta\beta$: Observed 2nd Order Weak Process.


$${}^Z A \Rightarrow {}^{Z+2} A + 2e^{-} + 2\bar{\nu}_e$$



Experimental Signature:

All the energy is shared between the 2 electrons: **monochromatic line** at the Q-value of the decay.

Examples $\Rightarrow \left\{ \begin{array}{ll} {}^{130}\text{Te} : & E_Q = 2528 \text{ keV} [4, 5] \\ {}^{76}\text{Ge} : & E_Q = 2039 \text{ keV} [6] \end{array} \right.$



Experimental Sensitivity $S^{0\nu}$:

Defined as the decay time corresponding to the minimum number of detectable events above background (B).

$$S^{0\nu} \propto \epsilon \cdot i.a. \cdot \sqrt{\frac{M \cdot T}{B \cdot \Gamma}}$$

Isotopic Abundance (points to i.a.)
 Detector Mass (points to M)
 Live time (points to T)
 Efficiency (points to ϵ)
 Background Counts (points to B)
 Energy Resolution (points to Γ)

Experiments can measure the decay rate $\Gamma_{1/2}^{0\nu}$


$$\Gamma_{1/2}^{0\nu} = [\tau_{1/2}^{0\nu}]^{-1} = G^{0\nu} |M^{0\nu}|^2 |\langle m_\nu \rangle|^2, \quad \left\{ \begin{array}{ll} G^{0\nu} & : \text{Phase Space Factor;} \\ |M^{0\nu}|^2 & : \text{Nuclear Matrix Elements;} \\ |\langle m_\nu \rangle|^2 & : \text{Neutrino Majorana Mass.} \end{array} \right.$$

where $|\langle m_\nu \rangle|^2 = |\sum_{i=1}^{n=3} m_i U_{ei}^2|$ is the **Effective Majorana Mass**

Observation of $0\nu\beta\beta$ would prove that

- ❶ lepton number is not conserved ($\Delta L = 2$);
- ❷ neutrinos are Massive Majorana Particles ($\nu = \bar{\nu}$).

It could also give informations on the absolute scale of neutrino masses and on the neutrino mass hierarchy (through $|\langle m_\nu \rangle|$)



We will discuss only selected next generation double-beta decay experiments that are approved and under construction:



- *COBRA*
- *CUORE*
- *EXO-200*
- *GERDA*
- *Kamiokande-Xe*
- *Majorana Demonstrator*



The idea




K. Zuber, Phys. Lett. B 519,1 (2001)

Use large amount of CdZnTe Semiconductor Detectors




- Source = detector
- Focus on ^{116}Cd
- Semiconductor (Good energy resolution, clean)
- Room temperature
- Modular design (Coincidences)
- Tracking/Pixelisation („Solid state TPC“)

Setup at Gran Sasso Lab



Current spectrum (black)

Background at 2810 keV about 5 cts/keV/kg/yr








Currently ongoing upgrade:

- 64 detectors (in hand)
- new DAQ
- Pulse shape information (done)
- Improvement on shielding
- new location at LNGS (former HdMo cabin)

The pixel option – Semiconductor tracker


Idea: Massive background reduction by particle identification



Particle identification works !!!

Preselected samples

256x256 pixels, 55 μ m



According to simulations particle identification due to pixels should reduce background by 3 orders of magnitude



CUORE

***Cryogenic Underground Observatory for
Rare Events***

***A search for neutrino-less double beta
decay of Te-130 in the Laboratori
Nazionali del Gran Sasso, Assergi, Italy***

***CUORE uses the bolometric technique
with TeO₂ single crystal bolometers.***

CUORE: the collaboration



Present Collaboration


64 European collaborators

32 US collaborators

5 Chinese collaborators

CUORE/Cuoricino Bolometer

TeO₂ Bolometer: Source = Detector



For $E = 1 \text{ MeV}$:

$$\Delta T = E/C \approx 0.1 \text{ mK}$$


signal size: 1 mV

Time constant:

$$\tau = C/G = 0.5 \text{ s}$$

Energy resolution (FWHM):
 $\sim 5\text{-}10 \text{ keV at } 2500 \text{ keV}$


Single pulse example



I bolometri di Cuoricino

Cristallo Assorbitore




L'assorbitore è un cristallo **5x5x5 cm³** di **TeO₂** il quale contiene il nucleo candidato **¹³⁰Te**





Sensore di temperatura

Il segnale termico è misurato tramite un **Termistore di Ge NTD**

$$R(T) = R_0 \exp \sqrt{\frac{T_0}{T}}$$



CUORICINO[10, 11, 12] (2003 - 2008) is still the **largest bolometric** experiment ever realized.



$5 \times 5 \times 5 \text{ cm}^3$
790 g



$3 \times 3 \times 6 \text{ cm}^3$
330 g


CURICINO: 62 TeO_2 Crystals

- It took data from April 2003 to June 2008;
- 11 modules of 4 TeO_2 crystals
 $5 \times 5 \times 5 \text{ cm}^3$ (790 g);
- 2 modules of 9 TeO_2 crystals
 $3 \times 3 \times 6 \text{ cm}^3$ (330 g);
 - 14 crystals of no-enriched ^{130}Te (34.2% [8])
 - 2 crystals enriched in ^{130}Te (75% [13]);
 - 2 crystals enriched in ^{128}Te (82.3%);

Total mass of TeO_2 : 40.7 kg

Total mass of ^{130}Te : 11.3 kg

Total exposure: 19.75 kg.y of ^{130}Te (big crystals alone: 15.8 kg.y of ^{130}Te)



NME References

- 1 R(QRPA): F. Simkovic *et al.*, PRC 77 (2008) [14];
- 2 pn(QRPA): O. Civitarese *et al.*, JoPCS 173 (2009) [15];
- 3 ISM: J. Menendez *et al.*, NPA 818 (2009) [16];
- 4 IBM: J. Barea and F. Iachello, PRC 79 (2009) [17];

Background: (0.153 ± 0.006) c/keV/kg/y [18]

Lower Limit, Half-life: $\tau_{1/2}^{0\nu}(^{130}\text{Te}) > 2.8 \cdot 10^{24}$ y (90% C. L.) [18]

Upper Limit, Majorana Mass: $m_{\nu_e} < (300 \div 710)$ meV [18]

CUORICINO


2003 - 2008



1 Tower
 $44 + 18 = 62$ crystals
 $\simeq 11$ kg of ^{130}Te
 $B = 0.153$ c/keV/kg/y

CUORE-0


2011 - 2014




1 Tower
52 crystals
 $\simeq 11$ kg of ^{130}Te
 $B = (0.05 \div 0.18)$ c/keV/kg/y

CUORE

2013 - 2018



19 Towers, 52 crystals each
988 crystals
 $\simeq 206$ kg of ^{130}Te
 $B = (0.01 \div 0.001)$ c/keV/kg/y



CUORE-0


- A single tower (52 TeO_2 crystals) realized with the same procedure of CUORE;
 - 1 crystals from the same production line;
 - 2 same copper and PTFE;
 - 3 CUORE-like copper surface cleaning;
 - 4 same assembly line;
- The tower will be installed in Hall A cryostat (the same as CUORICINO);
- Many aspects can be analyzed with CUORE-0;
 - 1 detector performances;
 - 2 radioactive background;

CUORE-0 will be assembled in the next months at LNGS and then put into operation before the end of the year.



CUORE (Cryogenic Underground Experiment for Rare Events)

8



- 998 TeO_2 crystals $5 \times 5 \times 5 \text{ cm}^3$ (750g), arranged in 19 towers;
- 741 kg of $\text{TeO}_2 \rightarrow 206 \text{ kg}$ of ^{130}Te
- Resolution 5 keV @ 2615 keV (FWHM)
- Background aim: 0.01-0.001 c/keV/kg/y

Main concepts:

- Stringent controls on radioactivity
- Heavy shielded (Roman lead)
- High efficiency in bkg rejection thanks to close packet geometry




The EXO Collaboration

K.Barry, M.Hughes, R.MacLellan, E.Niner,
A.Piepkke, K.Pushkin
Physics Dept, U. of Alabama, Tuscaloosa AL
P.Vogel
Physics Dept Caltech, Pasadena CA
M.Dixit, K.Graham, C.Green, C.Hagemann,
C.Hargrove, E.Rollin, D.Sinclair, V.Strickland
Carleton University, Ottawa, Canada
C.Benitez-Medina, S.Cook, W.Fairbank Jr., K.Hall,
B.Mong
Colorado State U., Fort Collins CO
M.Moe
Physics Dept UC Irvine, Irvine CA
D.Akimov, I.Alexandrov, A.Burenkov, M.Danilov,
A.Dolgolenko, A.Karelin, A.Kovalenko,
A.Kuchenko, V.Stekhanov, O.Zeldovich
ITEP Moscow, Russia
B.Aharmim, K.Donato, J.Farine, D.Hallman,
U.Wichoski
Laurentian U., Sudbury, Canada
P. Fierlinger, F.Roseau
Technical University of Munich


H.Breuer, C.Hall, L.Kaufman, D.Leonard,
S.Slutsky, Y-R.Yen
U. of Maryland, College Park MD
J.Cook, T.Daniels, K.Kumar, P.Morgan, A.Pocar, B.
Schmoll, C.Sterpka
U. of Massachusetts, Amherst MA
M.Auger, D.Franco, G.Giroux, R.Gornea, F.Juget,
G.Lutter, J-L.Vuilleumier, J-M.Vuilleumier
*Laboratory for High Energy Physics, Bern,
Switzerland*
N.Ackerman, M.Breidenbach, R.Conley,
W.Craddock, S.Herrin, J.Hodgson, D.Mackay,
A.Odian, C.Prescott, P.C.Rowson, K.Skarpaas,
M.Swift, K.Wamba, J.Wodin, L.Yang, S.Zalog
SLAC, Menlo Park CA
P.Barbeau, L.Bartoszek, L.Coleman, R.Cooper,
R.DeVoe, M.Dolinski, P.Fierlinger, B.Flatt,
G.Gratta, M.Green, T. Koffas, F.LePort,
M.Montero-Diez, R.Neilson, A.Reimer-Muller,
A.Rivas, K.O'Sullivan, K.Twelker
Physics Dept Stanford U., Stanford CA

EXO-200



- Ionization along track is collected on X-Y wires (at 60 degree angle)
- Scintillation light (175nm) is collected on Avalanche Photo Diodes (APDs)
- Energy reconstruction from combined ionization and scintillation signals yielding predicted $\Delta E/E = 1.4\%$ at $Q_{\beta\beta}$
- Position reconstructed from charge distribution and $t_{SCINT}-t_{ION}$; needed for Ba extraction in full scale EXO

EXO chamber "hugs" the fiducial volume very closely!




EXO-200 TPC basics

- The detector measures both the ionization electrons and the scintillation light to get best energy resolution.
 - In addition, the position of the decay is measured to get spatial distributions and (for later) the position of the Ba ion.
 - Info on event topology also important for background separation.
- The detector is a cylinder of ~40cm ID by ~40cm inner length.
- The cylinder is split by a cathode plane at the center so there are two symmetric drift regions. The cathode runs at negative HV.
 - Max HV is ~ 70kV (~3.5 kV/cm drift). Energy resolution improves with drift field, but there are arguments that separation of 1 vs 2 primary electrons might be better at lower fields.
 - field optimization is an important mission of EXO-200
- Readout "style":
 - Crossed wires, 100 μ m wires, 3mm pitch, ganged in groups of 3
 - 48ch x, 48ch y, total 96 ch per 1/2 detector
 - (Pad readout rejected because of high channel count)


EXO-200

Central HV plane
(photoetched
phosphor bronze)





~40cm
diameter

~35cm



The TPC vessel sits in a low background, copper cryostat filled with ultra-clean HFE7000 refrigerant/shielding fluid






TPC Insertion - 1/13/10

02/13/2010

Simon Slutsky APS April Meeting

11



Front shield
& Rn enclosure

Veto counter installed
and commissioned


Low background
data taking started




Two new ^{76}Ge Projects:



GERDA



Majorana



- 'Bare' ^{76}Ge array in liquid argon
- Shield: high-purity liquid Argon / H_2O
- Phase I: 18 kg (HdM/IGEX) / 15 kg nat.
- Phase II: add ~20 kg new enr. Detectors; total ~40 kg enriched detectors

- Array(s) of ^{76}Ge housed in high-purity electroformed copper cryostat
- Shield: electroformed copper / lead
- Initial phase: R&D demonstrator module: Total ~60 kg (30 kg enr.)

Physics goals: degenerate mass range

Technology: study of bgds. and exp. techniques


Lol

- open exchange of knowledge & technologies (e.g. MaGe MC)
- intention to merge for O(1 ton) exp. (inv. Hierarchy) selecting the best technologies tested in GERDA and Majorana

Characteristics of ^{76}Ge for $0\nu\beta\beta$ search

- **Favorable nuclear matrix element** $|M^{0\nu}|=3 - 9$
- **Reasonable slow $2\nu\beta\beta$ rate** ($T_{1/2} = 1.4 \times 10^{21}$ y) and **high $Q_{\beta\beta}$ value (2039 keV)**
- **Ge as source and detector**
- **Elemental Ge** maximizes the source-to-total mass ratio
- **Industrial techniques** and facilities available to enrich from 7% to ~86%
- **Intrinsic high-purity Ge diodes & HP-Ge detector technologies** well established
- **Excellent energy resolution:** FWHM ~3 keV at 2039 keV (0.16%)
- **Powerful signal identification & background rejection possible with novel detector concepts:**
 - time structure of charge signal (PSA) using **BEGe detectors**
 - granularity
 - **liquid argon scintillation as active veto system**
- **Best limits** on $0\nu\beta\beta$ - decay used Ge (IGEX & Heidelberg-Moscow)
 $T_{1/2} > 1.9 \times 10^{25}$ y (90%CL) [**& claim for evidence**]


Sum spectrum of the ^{76}Ge detectors Nr. 1,2,3,4,5



H.V. Klapdor-Kleingrothaus et al. NIM A522 (2004) 371-406, PLB 586 (2004) 198-212.

The Single Site Selected Spectrum of the ^{76}Ge detectors Nr. 2,3,4,5

HEIDELBERG-MOSCOW, 2004



Energy Range 100 - 3000 keV

H.V. Klapdor-Kleingrothaus et al.

$$T_{1/2}^{0\nu}({}^{76}\text{Ge}) = 2.23_{-0.31}^{+0.44} \times 10^{25} \text{ y}$$

H.V. Klapdor-Kleingrothaus and his colleagues reanalyzed the Heidelberg - Moscow Experiment data and concluded that the data does support the claim of first observation of zero-neutrino double-beta decay.

This has been highly controversial; however, it cannot be refuted nor confirmed except by experiment.



~ 100 members
19 institutions
6 countries



M. Agostini^a, M. Allardt^c, E. Andreotti^e, A.M. Bakalyarov^l, M. Balata^a,
I. Barabanov^j, M. Barnabe-Heider^f, L. Baudis^s, C. Bauer^f, N. Becerici-Schmid^m,
E. Bellotti^{g,h}, S. Belogurov^{k,j}, S.T. Belyaev^l, G. Benato^{o,p}, A. Bettini^{o,p}, L. Bezrukov^j,
T. Bruch^s, V. Brudanin^d, R. Brugnera^{o,p}, D. Budjasⁿ, A. Caldwell^m, C. Cattadori^{g,h},
F. Cossavella^m, E.V. Demidova^k, A. Denisov^j, S. Dinter^m, A. Domula^c, V. Egorov^d,
F. Faulstich^m, A. Ferella^s, K. Freund^r, F. Froborg^s, N. Frodyma^b, A. Gangapshv^j,
A. Garfagnini^{o,p}, S. Gazzana^{l,d}, P. Grabmayr^r, V. Gurentsov^j, K.N. Gusev^{l,d},
W. Hampel^f, A. Hegai^r, M. Heisel^f, S. Hemmer^{o,p}, G. Heusser^f, W. Hofmann^f,
M. Hult^e, L. Ianucci^a, L.V. Inzhechik^j, J. Janicskoⁿ, J. Jochum^r, M. Junker^a,
S. Kianovsky^j, I.V. Kirpichnikov^k, A. Kirsch^f, A. Klimenko^{d,j}, K-T. Knoepfle^f,
O. Kochetov^d, V.N. Kornoukhov^{k,j}, V. Kusminov^j, M. Laubenstein^a, V.I. Lebedev^l,
B. Lehnert^c, S. Lindemann^f, M. Lindner^f, X. Liu^q, A. Lubashevskiy^f,
B. Lubsandorzhiev^j, A.A. Machado^f, B. Majorovits^m, G. Marissens^e, G. Meierhofer^r,
I. Nemchenok^d, S. Nisi^a, C. O'Shaughnessy^m, L. Pandola^a, K. Pelczar^b, F. Potenza^a,
A. Pulliaⁱ, M. Reissfelder^f, S. Riboldiⁱ, F. Ritter^r, C. Sada^{o,p}, J. Schreiner^f,
U. Schwan^f, B. Schwingenheuer^f, S. Schönertⁿ, H. Seitz^m, M. Shirchenko^{l,d},
H. Simgen^f, A. Smolnikov^f, L. Stanco^p, F. Stelzer^m, H. Strecker^f, M. Tarka^s,
A.V. Tikhomirov^l, C.A. Ur^p, A.A. Vasenko^k, O. Volynets^m, M. Wojcik^b, E. Yanovich^j,
P. Zavarise^a, S.V. Zhukov^l, D. Zinatulina^d, F. Zoccaⁱ, K. Zuber^c, and G. Zuzel^b.

^a) INFN Laboratori Nazionali del Gran Sasso, LNGS, Assisi

^b) Institute of Physics, Jagellonian University, Cracow, Poland

^c) Institut für Kern- und Teilchenphysik, Technische Universität Dresden

^d) Joint Institute for Nuclear Research, Dubna, Russia

^e) Institute for Reference Materials and Measurements, Geel, Belgium

^f) Max Planck Institut für Kernphysik, Heidelberg, Germany

^g) Dipartimento di Fisica, Università Milano Bicocca, Milano, Italy

^h) INFN Milano Bicocca, Milano, Italy

ⁱ) Dipartimento di Fisica, Università degli Studi di Milano e INFN Milano

^j) Institute for Nuclear Research of the Russian Academy of Science

^k) Institute for Theoretical and Experimental Physics, Moscow, Russia

^l) Russian Research Center Kurchatov Institute, Moscow, Russia

^m) Max-Planck-Institut für Physik, München, Germany

ⁿ) Physik Department E15, Technische Universität München

^o) Dipartimento di Fisica dell'Università di Padova, Padova, Italy

^p) INFN Padova, Padova, Italy

^q) Shanghai Jiaotong University, Shanghai, China


^r) Physikalisches Institut, Eberhard Karls Universität Tübingen, Tübingen, Germany

^s) Physik Institut der Universität Zürich, Zürich, Switzerland



Background reduction:

Deep underground site for suppression of cosmic ray muons &
graded shielding against ambient radiation &
rigorous material selection &
signal analysis





Phase II Detectors:

“True-coaxial” natural detectors





mock-up



- 6-fold- ϕ segmented p-type
- 18-fold (6- ϕ ; 3-z) segmented n-type

18-fold segmented detector
(in standard cryostat)




© Jan Hattenbach

Inauguration of the GERDA experiment at the LNGS 9. November 2010





Phases and physics reach



required for 'background free' exp. with $\Delta E \sim 3.3$ keV (FWHM):

$O(10^{-3})$ $O(10^{-4})$ counts/(kg·y·keV)

Background requirement for GERDA/Majorana:

⇒ Background reduction by factor $10^2 - 10^3$ required w.r. to precursor exps.

⇒ Degenerate mass scale $O(10^2 \text{ kg} \cdot \text{y}) \Rightarrow$ Inverted mass scale $O(10^3 \text{ kg} \cdot \text{y})$



Conclusions & Outlook




- GERDA experimental **installations completed** successfully; cryogenic and auxiliary systems operate very stable
- Detector **commissioning** with **non-enriched detectors** started summer 2010 and is still ongoing
- **Initial count rate** (run 1-3,5) **dominated by ^{42}K (^{42}Ar progenitor)** due to concentration of ^{42}K close to the detectors by E-field of diodes $\Rightarrow$ **field-free configuration**
- **12 commissioning runs** with different detectors, read-out schemes, E-field configurations completed successfully
- Background with **non-enriched detectors currently at 0.05 cts/(keV kg year)**. Goal for Phase I: 0.01 cts/(keV kg year).
- Deployment of first string(s) with **enriched detectors**
Phase I soon to study background with enriched detectors
 $\Rightarrow$ start of Phase I physics run



Conclusions & Outlook (cont.)

- Thick-window **p-type BEGe** detectors for **Phase II**
- Powerful **particle ID** and **background discrimination** by pulse shape analysis: MSE/SSE, p+ contact (α) and n+ (β) surface events
- Full **production chain tested** for BEGe Phase II detectors
- 37.5 kg of 86% ^{enr}**Ge** (in form of GeO₂) successfully transformed to **35.4 kg** (94%) of 6N
- Crystal **pulling** and **detector production** under preparation
- **Liquid argon instrumentation** shown in **GERDA-LArGe** test stand to be a powerful method to discriminate backgrounds: **implementation in GERDA if needed**



The MAJORANA DEMONSTRATOR





- The MAJORANA experiment is a US-led effort that will be deployed at the Sanford lab in South Dakota
- The design focuses on the use of high purity material and will use a Pb/Cu shield
- The experiment will be run in phases and will culminate with the use of 60 kg of Ge (30kg of which will be enriched)

[illegible]


-




Point contact Ge detector




Point contact Ge detector allow multi-site events to be identified-invented by Paul Luke (LBNL)



Pulse-shape discrimination



Segmentation






Fig. 1. The left figure shows charge-drift paths (dark) and isochrone lines (light) of a point-contact detector. Two events on the same isochrone appear as a single event. The color scale indicates the drift velocity of holes (in mm/ns). The right figure shows the charge pulse (top) and current pulse (lower) of a multisite event, from a gamma ray.







Fig. 2. Artist's conception of the DEMONSTRATOR shield (left panel), and the result of applying pulse-shape analysis on a spectrum from a calibration source of ^{232}Th irradiating a BEGe detector (right panel).

Electroformed copper




- The use of electroformed copper removes impurities and significantly reduces backgrounds; goal is $0.3 \mu\text{Bq } ^{232}\text{Th}$ and $^{238}\text{U/kg Cu}$ ($\sim 0.08 \times 10^{-12} \text{ g/gCu}$)
- The copper is electroformed on steel mandrels that have the same diameter as the cryostats
- A total of 16 baths will be deployed underground. The material from the electroformed cylinders will be machined into various components

Thermosiphon cooling for coldplate





- Solid Copper cold finger temp rise $\sim 17\text{K}$
- Thermosiphon temp rise $< 5\text{K}$
- Better scaling to 1-tonne cryostat
- Shorter e-forming production time




Detector Simulation

- The MAJORANA and GERDA collaborations have worked on the MaGe Monte Carlo simulation package (using Geant4 and ROOT)
- O(5) publications
- Pulse shape calculations also implemented
- Simulation and Analysis task lead is at LBNL





Estimated Schedules



Towards a 1 tonne Ge experiment


Sensitivity depends on energy resolution, background rate and exposure:

$$\langle m_{\beta\beta} \rangle \propto \left(\frac{b\Delta E}{Mt_{\text{live}}} \right)^{\frac{1}{4}}$$

background rate $\rightarrow b$ energy resolution $\rightarrow \Delta E$

active mass $\rightarrow M$ live-time $\rightarrow t_{\text{live}}$

Half life (years)	~Signal (cnts/ton-year)	~Neutrino mass scale (meV)
10^{25}	530	400
5×10^{26}	10	100
5×10^{27}	1	40
$> 10^{29}$	< 0.05	< 10





A 1 tonne experiment is well suited for exploring mass scales down to the atmospheric neutrino oscillation mass-squared difference.

Perspectives for $2\beta 0\nu$ Search at KamLAND




Yuri Efremenko: UTK, IPMU
MEDEX'11: June, Prague 2011
(On behalf of the KamLAND-Zen collaboration)

KAMLAND



The neutron slows down in the scintillator and is captured in a Gd nucleus. This results in capture-gamma rays delayed in time from the



Xenon

Noble gas

Dissolvable in Liquid Scintillator

Density 5.894 g/l

Melting point -111.7 °C

Boiling point -108.2 °C

World production ~27 t/y

Applications: Illumination, Anesthesia, Particle detectors, Ion thrusters



¹³⁶Xe

Natural Abundance: 8.9%

Q value: 2476 keV

$T_{1/2}$ $2\beta 2\nu$ >10²² y.

$T_{1/2}$ $2\beta 0\nu$ for 50 meV ~3.0÷4.4·10²⁶ y.

KamLAND Zen

Mini balloon

Radius:	1.58m
Material:	nylon-6
Thickness	25 μ m
Density	1.14 g/cm ³
Weight	900g

Xe loaded liquid scintillator


91.7% enriched ¹³⁶Xe 400kg (3.0wt%)

Composition ratio

Decane	82.3 %
Pseudocumene	17.7 %
PPO	2.7g/l

Density : same as KamLAND liquid
scintillator (0.777 g/cm³)

Light yield : same as KamLAND liquid
scintillator




Energy Resolution: 6.8%/ $\sqrt{E(\text{MeV})}$
Vertex Resolution: 12.5cm/ $\sqrt{E(\text{MeV})}$

Mini Balloon Status


By Shirai-san

*Super clean room is ready !
(at Nishizawa center of Tohoku Univ.)*


Class 1 (=1 particle(>0.1 μ m) /feet³)




*Welding method and conditions
has been established.*




*Final checks and preparation of
the balloon construction and
deployment are ongoing.*



*Designing the
monitor cameras
and lights in the
deployment are
finished and will
be ordered soon.*



KamLAND-Zen Sensitivity



Present Status and Plans

Xe handling system is in place

370 kg of ^{136}Xe are in the mine (50 kg are coming soon)

Xe LS system has been assembled

Mini Balloon is being made

Final Calibration for Reactor phase is ongoing

Scintillator mixing – July

Chimney modification – July

Mini Balloon deployment and filing – August

Data Taking from September

First result later this year

Other Experiments

Experiment	Isotope	Mass	Technique	Present Status	Location
CANDLES	^{48}Ca	0.35 kg	CaF_2 scint. crystals	Prototype - 2009	Kamioka
CARVEL	^{48}Ca	1 ton	CaF_2 scint. crystals	Development	Solotvina
COBRA	^{116}Cd	183 kg	^{enr}Cd CZT semicond. det.	Prototype	Gran Sasso
CUORICINO	^{130}Te	11 kg	TeO_2 bolometers	Complete - 2008	Gran Sasso
CUORE	^{130}Te	200 kg	TeO_2 bolometers	Construction - 2012	Gran Sasso
DCBA	^{150}Nd	20 kg	^{enr}Nd foils and tracking	Development	Kamioka
EXO-200	^{136}Xe	160 kg	Liq. ^{enr}Xe TPC/scint.	Construction - 2009	WIPP
EXO	^{136}Xe	1-10 t	Liq. ^{enr}Xe TPC/scint.	Proposal	DUSEL
GEM	^{76}Ge	1 ton	^{enr}Ge det. in liq. nitrogen	Inactive	
GENIUS	^{76}Ge	1 ton	^{enr}Ge det. in liq. nitrogen	Inactive	
GERDA	^{76}Ge	≈ 35 kg	^{enr}Ge semicond. det.	Construction - 2009	Gran Sasso
GSO	^{160}Gd	2 ton	$\text{Gd}_2\text{SiO}_5:\text{Ce}$ crys. scint. in liq. scint.	Development	
KamLAND	^{136}Xe	200 kg	^{enr}Xe dissolved in liq. scint.	Construction - 2012?	Kamioka
MAJORANA	^{76}Ge	26 kg	^{enr}Ge semicond. det.	Construction - 2011	SUL
MOON	^{100}Mo	1 t	^{enr}Mo foils/scint.	Development	
NEXT	^{136}Xe	80 kg	gas TPC	Development	Canfranc
SNO+	^{150}Nd	55 kg	Nd loaded liq. scint.	Construction - 2011	SNOLab
SuperNEMO	^{82}Se	100 kg	^{enr}Se foils/tracking	Proposal	Frejus
Xe	^{136}Xe	1.56 t	^{enr}Xe in liq. scint.	Development	
XMASS	^{136}Xe	10 ton	liquid Xe	Inactive for $\beta\beta$	Kamioka
HPXe	^{136}Xe	tons	High Pressure Xe gas	Development	

FINAL CONCLUSION

The Next Generation neutrino less double-beta decay experiments should:

- 1. Determine the Validity of the KKDK Claim of Discovery***
- 2. If Not, then Probe Neutrino Mass Scales of 50-100 milli-eV***

For a more detailed discussion see:

F.T. AVIGNONE III, S.R. ELLIOTT and J. ENGEL

“Double beta decay, Majorana neutrinos, and neutrino mass,”

REVIEWS OF MODERN PHYSICS VOL. 80, APRIL-JUNE 2008.